

cedure was done, however, the small amount of pituitary tissue in the otherwise empty sella had basophilic hyperplasia and no adenoma. In addition, partial pituitary hypofunction is suggested by plasma gonadotropin levels that were inappropriately low for her postmenopausal state. Thyroid function was normal. Several management issues arose during the course of this patient's illness that deserve comment. Selective venous catheterization would have been helpful in establishing the pituitary as the source of inappropriate ACTH production. In a case of pituitary-dependent Cushing's and an empty sella, it is probably still warranted to surgically explore the pituitary fossa for a possible adenoma. Bilateral adrenalectomy remains, however, a reasonable treatment alternative.

The question of whether this patient had Cushing's disease or Cushing's syndrome on another basis cannot ultimately be settled without a postmortem examination. The weight of the evidence suggests a pituitary basis. The factors supporting a primary pituitary aberration are the inappropriate rise in the ACTH level (though still within the normal range) in the presence of extreme hypercortisolism and its partial suppressibility. Other possible causes for Cushing's syndrome appear to be reasonably excluded. Documentation by CT scan of enlargement of both adrenal glands rules out a unilateral adrenocortical adenoma as the source of this patient's hypercortisolism. An ectopic source for ACTH is also unlikely for several reasons. The patient's long clinical course (six years), negative findings on a chest x-ray film, normal 24-hour urinary excretion of 5-hydroxyindoleacetic acid and partially suppressible hypercortisolism are inconsistent with an ectopic source for ACTH. Furthermore, the ACTH level in this paraneoplastic syndrome tends to be much higher than occurred in this patient. The failure of high-dose dexamethasone to suppress urinary 17-hydroxycorticosteroid excretion by at least 50% may argue against pituitary-dependent Cushing's disease, but 15% to 30% of these patients fail to suppress to this degree.¹⁴

This case also has potentially interesting implications concerning the natural history of Cushing's disease. If this patient's primary defect was at the level of the hypothalamus resulting in inappropriate stimulation of the pituitary, this would have caused development of a pituitary adenoma, ACTH hypersecretion and cushingoid features. Infarction of the adenoma may have then occurred, resulting in an empty sella. Of other possible causes of an empty sella, a developmentally incomplete sellar diaphragm cannot be completely excluded, whereas involution of a previously hypertrophied and overstimulated pituitary due to end-organ failure appears highly unlikely.

REFERENCES

1. Weisberg LA, Zimmerman EA, Frantz AG: Diagnosis and evaluation of patients with an enlarged sella turcica. *Am J Med* 1976 Nov; 61:590-596
2. Ganguly A, Stanchfield JB, Roberts TS, et al: Cushing's syndrome in a patient with an empty sella turcica and a microadenoma of the adenohypophysis. *Am J Med* 1976 Feb; 60:306-309
3. Gutin PH, Cusard WG Jr, Wilson CB: Cushing's disease with pituitary apoplexy leading to hypopituitarism, empty sella, and spontaneous fracture of the dorsum sellae. *J Neurosurg* 1979 Dec; 51:866-869
4. Smith JS, Kohler PC, Helminiak R, et al: Intermittent Cushing's syndrome with an empty sella turcica. *Arch Intern Med* 1982; 142:2185-2187
5. Sutton TJ, Vezina JL: Co-existing pituitary adenoma and intrasellar arachnoid invagination. *Am J Roentgenol Radium Ther Nucl Med* 1974 Nov; 122:508-510
6. Domingue JN, Wing SD, Wilson CB: Coexisting pituitary adenomas and partially empty sellas. *J Neurosurg* 1978; 48:23-28
7. Bar RS, Mazzaferri EL, Malarkey WB: Primary empty sella, galactorrhea, hyperprolactinemia and renal tubular acidosis. *Am J Med* 1975 Dec; 59:863-866
8. Hsu TH, Shapiro JR, Tyson JE, et al: Hyperprolactinemia associated with empty sella syndrome. *JAMA* 1976 May 3; 235:2002-2004
9. Bryner JR, Greenblatt RB: Primary empty sella syndrome with elevated serum prolactin. *Obstet Gynecol* 1977 Sep; 50:375-380
10. Login I, Santen RJ: Empty sella syndrome—Sequelae of the spontaneous remission of acromegaly. *Arch Intern Med* 1975 Nov; 135:1519-1521
11. Sindler BH, Griffing GT, Melby JC: Primary empty sella syndrome, ACTH hypersecretion and normal adrenocortical function—Report of two cases. *Am J Med* 1982; 73:449-454
12. Sindler BH, Melby JC: Primary empty sella syndrome, ACTH hypersecretion, and normal cortisol dynamics: Evidence for large molecular weight bioinactive ACTH. *The Endocrine Society, Programs and Abstracts, 64th Annual Meeting, San Francisco, 1982, p 149*
13. Jara-Albarran A, Bayart J, Caballero A, et al: Probable pituitary adenoma with adrenocorticotropin hypersecretion (corticotropinoma) secondary to Addison's disease. *J Clin Endocrinol Metabol* 1979 Aug; 49:236-241
14. Aron DC, Tyrell JB, Fitzgerald PA, et al: Cushing's syndrome: Problems in diagnosis. *Medicine (Baltimore)* 1981 Jan; 60:25-35

Homonymous Hemianopia Due to Cerebral Air Embolism From Central Venous Catheters

PATRICK J. KEARNS, JR, MD
ANTHONY A. HAULK, MD
THOMAS W. McDONALD, MD
San Jose, California

PERCUTANEOUS PLACEMENT of a central venous catheter has become a routine bedside procedure. Immediate mechanical complications include direct trauma to nerves, arteries and pleura and embolization of air or catheter fragments. Delayed complications are primarily infectious and thrombotic in nature.^{1,2} The following reports of cases illustrate an unusual complication of central venous catheters—homonymous hemianopia due to cerebral air embolism—and highlight the need to implement precautions to prevent this complication.

Reports of Cases

CASE 1. The patient, a 60-year-old man, was admitted for control of malignant hypertension and unstable angina. Evaluation showed diffuse arteriosclerosis, which included carotid and renal artery disease as well as 70% occlusion of his left anterior descending and right coronary arteries. The patient underwent right carotid endarterectomy in April 1981 and coronary-artery-bypass graft operation three weeks later. Both procedures were uneventful. Five days after the cardiac operation the patient was transferred to the ward with a left subclavian venous catheter in place.

(Kearns PJ Jr, Haulk AA, McDonald TW: Homonymous hemianopia due to cerebral air embolism from central venous catheters. *West J Med* 1984 Apr; 140:615-617)

From the Department of Medicine, Santa Clara Valley Medical Center, San Jose, California.

Submitted, revised, June 22, 1983.

Reprint requests to Patrick J. Kearns, Jr, MD, Dept of Medicine, Santa Clara Valley Medical Center, 751 S Bascom Ave, San Jose, CA 95128.

Two hours later, the patient unwittingly disconnected the catheter tubing to walk to the bathroom. He was found standing in the doorway with the central venous catheter disconnected, complaining of shortness of breath and back, shoulder and jaw pain. On physical examination, the patient had a heart rate of 120 per minute, a respiratory rate of 30 and a blood pressure of 200/170 mm of mercury. No blood was noted to escape from the disconnected catheter. He had a left homonymous hemianopia and diminished sensation of the left side of his body; both findings were new. His supine central venous pressure measured 4 cm of water, which remained stable for the next 24 hours. The patient was placed at bed rest and given high-flow oxygen therapy. His blood pressure gradually fell to 130/80 mm of mercury during the first hour and his respirations returned to 20 per minute.

A diagnosis of air embolism from the disconnected catheter was considered. A computed tomogram (CT) of the head done 12 hours after the initial symptoms showed a circumscribed 4-mm air-density lesion (about -400 Hounsfield units) in the right calcarine cortex consistent with an air embolus in the right posterior cerebral artery (Figure 1). This finding correlated well with the patient's symptoms of left homonymous hemianopia and left-sided sensory abnormalities. The patient's sensory deficits totally resolved during the ensu-




Figure 1.—Computed tomogram of the patient in case 1 obtained 12 hours after the onset of symptoms, showing an air-density lesion (arrow). A repeat scan 5 days later showed resolution of the air-density lesion with a residual area of low attenuation.

ing five days, and the visual field defect improved slightly. A repeat CT scan five days after the initial study showed complete resolution of the air embolus, with a residual area of low attenuation in the right occipital cortex consistent with cerebral infarction. Review of the findings of the physical examination, cardiac catheterization and echocardiogram showed no evidence of ventricular or atrial septal defect or a patent foramen ovale. He was discharged and readmitted eight months later for aortofemoral bypass with renal artery reconstruction. At that time he had only a residual inferior left quadrantanopia.

CASE 2. The patient, a 63-year-old man with diabetes mellitus and coronary artery disease, was admitted because of bacteremia, cellulitis and staphylococcal abscesses of the left lower extremity. The patient had poor peripheral venous access and required a central venous catheter for intravenous administration of antibiotic therapy. Two days after placement of the catheter, he inadvertently removed the catheter despite stabilization by a suture ligature. A sterile dressing was applied. Within an hour the patient noted decreased vision in his left eye, but he did not report this symptom at that time. A second central venous catheter was placed without incident, and antibiotic therapy was continued. Five days after placement of the second catheter, the patient complained of persistent left eye "blindness" and stated that the onset had occurred at the time of the dislodgement of the first central venous catheter. On physical examination a left homonymous hemianopia was found. A CT scan showed an area of low attenuation in the occipital cortical region consistent with cerebral infarction. Digital subtraction angiography showed no evidence of carotid artery disease. Neither echocardiogram nor cardiac catheterization showed evidence of a mural thrombus or valvular vegetation. A two-week course of antibiotics was completed and the infection resolved. The left homonymous hemianopia was unchanged when the patient was seen a year following the cerebral infarction. The CT scan has remained unchanged.

Discussion

The case of a patient who had a cerebral air embolus from a disconnected central venous catheter with a transient homonymous hemianopia was reported by Menkin and Schwartzman.³ Postulated mechanisms for the neurologic symptoms following air embolism include obstruction of the outflow tract of the right ventricle with consequent decrease in cardiac output, intense cerebral vasospasm induced by air emboli passing through intracranial vessels and actual occlusion of cerebral vessels by trapped air emboli.^{4,5}

The findings of the CT scan in the first patient may be the first clear clinicoanatomic correlation between neurologic symptoms and an air embolus in the brain. In this patient the underlying pathophysiologic feature was obstruction to cerebral blood flow by the air embolus, which caused ischemia and infarction. The timing of the CT scan showed that clinically significant air

TABLE 1.—Accumulated Experience of Cerebral Air Emboli

Study, Year	Patients N*	Catheter Associated N	Neurologic Symptoms (N)	Mortality %
Green and Nemir, 1971 ¹¹	1	1	Blindness (1), paresis	...
Hoshal and Fink, 1969 ¹²	2	2	Semicoma (1)	...
Ponsky, 1971 ⁹	1	1	Coma (1), seizures	100
Menkin and Schwartzman, 1973-1974 ³	5	1	Hemianopia (1)	40
Coppa et al, 1976-1980 ¹³	14	14	Coma, seizure (9), hemiplegia	30
Grace, 1977 ¹⁴	2	2	Hemiparesis (2), coma	...
Peters and Armstrong, 1978 ⁵	2	2	Monoplegia (1)	50
Michel et al, 1982 ¹⁵	1	1	Coma (1)	100
Eisenhauer et al, 1982 ¹⁶	3	3	Seizure (2)	...
TOTALS	31	19		

*Total number of patients in cited report who had presumed air emboli.

emboli can persist up to 12 hours and are resorbed within five days.

The mechanism by which venous air emboli are delivered to the systemic circulation is controversial. The previously postulated mechanisms presumed obstruction of the outflow tract of the pulmonary artery by air embolism with an increase in the pulmonary artery pressure transmitted to the right atrium. There is subsequent shunting of blood through an existing anatomic defect, such as a patent foramen ovale.⁶ Alternative explanations must now be considered because the first patient had a low central venous pressure during the entire period of observation. In addition, neither an atrial septal defect nor a patent foramen ovale could be found. Intrapulmonary, physiologic right-to-left shunts have been found in dogs and in humans.⁷ These are presumed to be patent and functional and are of sufficient size to allow passage of air emboli. Butler and Hills showed in dogs that aminophylline and vasodilators "open up" the pulmonary venous-arterial barrier and allow passage of air emboli to the arterial circulation.⁸ The appearance of systemic air emboli with normal right-sided pressures in case 1 supports the significance of intrapulmonary shunts in the pathophysiology of the cerebral air embolus.

The neurologic symptoms of these two patients were assumed initially to be due to malignant hypertension, atherosclerotic disease or an idiopathic cause. Because of a high index of suspicion for cerebral air embolism, this diagnosis was confirmed in the first patient, and all other possible causes were excluded in the second patient. In all, 19 cases with the clinical diagnosis of cerebral air embolism due to central venous catheters have been reported, 15 within the past five years (Table 1). Because of the increasing use of central venous catheters for central pressure monitoring, for alimenting, in placing temporary pacemakers and in subclavian catheter dialysis, the incidence of air emboli with neurologic complications is likely to increase. Because these two patients were standing during their episodes, the air embolus followed a predictable route and traveled to the right posterior circulation. As noted by Ohkuda and co-workers,⁹ venous air emboli usually do

not cause symptoms, possibly because in supine patients the embolus can travel to neurologically silent areas. Because of this and the misconception that significant air emboli cause cardiovascular collapse, many air emboli may go unnoticed.

Because of the unknown incidence of this complication and the increasing use of the procedure, the following precautions are recommended for all central venous catheter placements:

- Follow proper insertion technique¹⁰;
- Use Luer-lock connections to the catheter to avoid disconnection;
- Place suture-ligatures to prevent movement of the catheter, best accomplished with catheters equipped with ligature eyelets.

If these simple precautions are followed, cerebral air embolism should be reduced. When the possibility of air embolism exists, the use of computed tomography for the early diagnosis of cerebral air embolism should expedite the diagnosis and institution of definitive therapy.

REFERENCES

1. Bernard RW, Stahl WM: Subclavian vein catheterization: A prospective study. *Ann Surg* 1971 Feb; 137:184-200
2. Johnson CL, Lazarchick JL, Lynn HB: Subclavian venipuncture: Preventable complications—Report of two cases. *Mayo Clin Proc* 1970 Oct; 45:712-719
3. Menkin M, Schwartzman RJ: Cerebral air embolism. *Arch Neurol* 1977 Mar; 34:168-170
4. Paskin DL, Hoffman WS, Tuddenham WJ: A new complication of subclavian catheterization. *Ann Surg* 1974 Mar; 179:266-268
5. Peters JL, Armstrong R: Air embolism occurring as a complication of central venous catheterization. *Ann Surg* 1978 Apr; 187:375-378
6. Ponsky JL: Paradoxical cerebral air embolism. *N Engl J Med* 1971 Apr 29; 284:985
7. Weibel ER: Design and structure of the human lung, chap 15, *In* Fishman AP (Ed): *Pulmonary Diseases and Disorders—Vol 1*. New York, McGraw-Hill, 1980, pp 224-271
8. Butler BD, Hills BA: The lung as a filter for microbubbles. *J Appl Physiol* 1979 Sep; 47:537-543
9. Ohkuda K, Nakahara K, Binder A, et al: Venous air emboli in sheep: Reversible increase in lung microvascular permeability. *J Appl Physiol* 1981 Oct; 51:887-894
10. Rosen M, Latio IP, Shang NGW: Percutaneous central venous cannulation. *Br Med J* 1980 Aug; 2:372-375
11. Green HL, Nemir P Jr: Air embolism as a complication during parenteral alimentation. *Am J Surg* 1971 May; 121:614-616
12. Hoshal VL, Fink GH: The subclavian catheter (Letter). *N Engl J Med* 1969 Dec 18; 281:1425
13. Coppa GF, Gouge TH, Hofstetter SR: Air embolism: A lethal but preventable complication of subclavian vein catheterization. *JPEN* 1981 Mar-Apr; 5:166-168
14. Grace DM: Air embolism with neurologic complications: A potential hazard of central venous catheters. *Can J Surg* 1977 Jan; 20:51-53
15. Michel L, Poskanzer DC, McKusick KA, et al: Fatal paradoxical air embolism to the brain: Complication of central venous catheterization. *JPEN* 1983 Jan-Feb; 6:68-70
16. Eisenhauer ED, Derveloy RJ, Hastings PR: Prospective evaluation of central venous pressure (CVP) catheters in a large city-county hospital. *Ann Surg* 1982 Nov; 196:560-564